

# Robustness analysis with the Alkire-Foster measures

Gastón Yalonetzky

OPHI-HDCA Summer School, Delft, 24 August - 3 September 2011.

We are grateful to the World Bank, two anonymous donors and OPHI for financial support

# Table of contents

Introduction

Stochastic dominance conditions for H and M0

Other methods of rank robustness

# Introduction: Robustness versus Dominance

- ▶ In general, robustness analysis assesses the sensitivity of classifications generated by an index, when the values of the index's key parameters change.

# Introduction: Robustness versus Dominance

- ▶ In general, robustness analysis assesses the sensitivity of classifications generated by an index, when the values of the index's key parameters change.
- ▶ Stochastic dominance conditions provide an extreme form of robustness: when they are fulfilled, a comparison is robust to a broad range of parameter values.

# Introduction: Robustness versus Dominance

- ▶ In general, robustness analysis assesses the sensitivity of classifications generated by an index, when the values of the index's key parameters change.
- ▶ Stochastic dominance conditions provide an extreme form of robustness: when they are fulfilled, a comparison is robust to a broad range of parameter values.

# Robustness of rankings generated by the AF measures (including the MPI)

The positions generated by AF measures may be sensitive to changes in the measure's key parameters, specifically:

## Robustness of rankings generated by the AF measures (including the MPI)

The positions generated by AF measures may be sensitive to changes in the measure's key parameters, specifically:

1. The poverty lines of each variable (i.e. the "first" cut-off):  $z_d$ .

## Robustness of rankings generated by the AF measures (including the MPI)

The positions generated by AF measures may be sensitive to changes in the measure's key parameters, specifically:

1. The poverty lines of each variable (i.e. the "first" cut-off):  $z_d$ .
2. The weights on each variable/dimension:  $w_d$ .



## Robustness of rankings generated by the AF measures (including the MPI)

The positions generated by AF measures may be sensitive to changes in the measure's key parameters, specifically:

1. The poverty lines of each variable (i.e. the "first" cut-off):  $z_d$ .
2. The weights on each variable/dimension:  $w_d$ .
3. The threshold that the weighted sum of deprivations need to surpass in order to identify someone as (multidimensionally) poor (i.e. the "second" cut-off):  $k$ .

# Statistical tools for the evaluation of robustness of rankings for the AF measures

There are, in principle, two approaches to evaluating the robustness of AF measures to changes in their key parameters:

# Statistical tools for the evaluation of robustness of rankings for the AF measures

There are, in principle, two approaches to evaluating the robustness of AF measures to changes in their key parameters:

1. To "fix" all parameter values and check the sensitivity to changes in a set of parameters.

# Statistical tools for the evaluation of robustness of rankings for the AF measures

There are, in principle, two approaches to evaluating the robustness of AF measures to changes in their key parameters:

1. To "fix" all parameter values and check the sensitivity to changes in a set of parameters.
  - ▶ Example: Batana (2008) fixes weights and poverty lines and checks the sensitivity of the rankings to changes in  $k$ .

# Statistical tools for the evaluation of robustness of rankings for the AF measures

There are, in principle, two approaches to evaluating the robustness of AF measures to changes in their key parameters:

1. To "fix" all parameter values and check the sensitivity to changes in a set of parameters.
  - ▶ Example: Batana (2008) fixes weights and poverty lines and checks the sensitivity of the rankings to changes in  $k$ .
  - ▶ Another one: Alkire y Foster (2009) y Lasso de la Vega (2009) derive dominance conditions over  $k$  keeping weights and poverty lines fixed.

# Statistical tools for the evaluation of robustness of rankings for the AF measures

There are, in principle, two approaches to evaluating the robustness of AF measures to changes in their key parameters:

1. To "fix" all parameter values and check the sensitivity to changes in a set of parameters.
  - ▶ Example: Batana (2008) fixes weights and poverty lines and checks the sensitivity of the rankings to changes in  $k$ .
  - ▶ Another one: Alkire y Foster (2009) y Lasso de la Vega (2009) derive dominance conditions over  $k$  keeping weights and poverty lines fixed.
2. To derive conditions under which an ordering is robust for all lines, weights and multidimensional cut-offs.

# Statistical tools for the evaluation of robustness of rankings for the AF measures

In this presentation we are going to review:

# Statistical tools for the evaluation of robustness of rankings for the AF measures

In this presentation we are going to review:

- ▶ Stochastic dominance conditions for  $H$  and  $M^0$  that ensure robustness for all multidimensional thresholds, weights and lines.



# Statistical tools for the evaluation of robustness of rankings for the AF measures

In this presentation we are going to review:

- ▶ Stochastic dominance conditions for  $H$  and  $M^0$  that ensure robustness for all multidimensional thresholds, weights and lines.
- ▶ Some basic robustness tests (applied to weights).

## The counting vector: key ingredient for the dominance conditions of H and M0

For person  $n$  we define  $D = c_n$ , where:

## The counting vector: key ingredient for the dominance conditions of H and M0

For person  $n$  we define  $D = c_n$ , where:

$$c_n = \sum_{d=1}^D w_d I(x_{id} \leq z_d)$$


## The counting vector: key ingredient for the dominance conditions of H and M0

For person  $n$  we define  $D - c_n$ , where:

$$c_n = \sum_{d=1}^D w_d I(x_{id} \leq z_d)$$

We then consider a distribution of deprivations,  $D - c$ , in the population, with values ranging from 0 (poor in every variable) to  $D$  (non-poor in every variable). A typical cumulative distribution is:

## A typical cumulative distribution of D-c



## The dominance condition over $k$

The key results are the following:


## The dominance condition over k

The key results are the following:

$$F^A(D-c) \leq F^B(D-c) \forall (D-c) \in [0, D] \leftrightarrow H^A \leq H^B \forall (D-c) \in [0, D]$$


$$H^A \leq H^B \forall (D-c) \in [0, D] \rightarrow M^A \leq M^B \forall (D-c) \in [0, D]$$

# The key result in one graph: I





## The key result in one graph: II



## Proof, as explained by Alkire and Foster

Notice that  $M0$  can be expressed in terms of  $H$  the following way:

$$M0(k) = \frac{1}{D} [H(D)D + \sum_{j=k}^{D-1} j[H(j) - H(j+1)]]$$

## Proof, as explained by Alkire and Foster

Notice that  $M0$  can be expressed in terms of  $H$  the following way:

$$M0(k) = \frac{1}{D} [H(D)D + \sum_{j=k}^{D-1} j[H(j) - H(j+1)]]$$

Simplifying that expression yields the following result:

$$M0(k) = \frac{1}{D} \left[ \sum_{j=k+1}^D H(j) + kH(k) \right]$$

## Proof, as explained by Alkire and Foster

Notice that  $M0$  can be expressed in terms of  $H$  the following way:

$$M0(k) = \frac{1}{D} [H(D)D + \sum_{j=k}^{D-1} j[H(j) - H(j+1)]]$$

Simplifying that expression yields the following result:

$$M0(k) = \frac{1}{D} \left[ \sum_{j=k+1}^D H(j) + kH(k) \right]$$

Therefore:  $H^A(k) \leq H^B(k) \forall k \in [1, D] \rightarrow M^A(k) \leq M^B(k) \forall k \in [1, D]$

## More dominance results: incorporating weights and poverty lines

We just saw that  $M^A(k) \leq M^B(k) \forall k \in [1, D]$  holds if:  
 $H^A(k) \leq H^B(k) \forall k \in [1, D]$

## More dominance results: incorporating weights and poverty lines

We just saw that  $M^A(k) \leq M^B(k) \forall k \in [1, D]$  holds if:  
 $H^A(k) \leq H^B(k) \forall k \in [1, D]$

Are there conditions under which  $H^A(k) \leq H^B(k) \forall k \in [1, D]$  for any vector of weights and poverty lines?

## More dominance results: incorporating weights and poverty lines

We just saw that  $M^A(k) \leq M^B(k) \forall k \in [1, D]$  holds if:  
 $H^A(k) \leq H^B(k) \forall k \in [1, D]$

Are there conditions under which  $H^A(k) \leq H^B(k) \forall k \in [1, D]$  for any vector of weights and poverty lines?

Yes, but so far we know of their existence only on two restrictive situations:

## More dominance results: incorporating weights and poverty lines

We just saw that  $M^A(k) \leq M^B(k) \forall k \in [1, D]$  holds if:  
 $H^A(k) \leq H^B(k) \forall k \in [1, D]$

Are there conditions under which  $H^A(k) \leq H^B(k) \forall k \in [1, D]$  for any vector of weights and poverty lines?

Yes, but so far we know of their existence only on two restrictive situations:

- ▶ When there are only two variables.



## More dominance results: incorporating weights and poverty lines

We just saw that  $M^A(k) \leq M^B(k) \forall k \in [1, D]$  holds if:  
 $H^A(k) \leq H^B(k) \forall k \in [1, D]$

Are there conditions under which  $H^A(k) \leq H^B(k) \forall k \in [1, D]$  for any vector of weights and poverty lines?

Yes, but so far we know of their existence only on two restrictive situations:

- ▶ When there are only two variables.
- ▶ For any number of variables under extreme identification approaches (union and intersection).

## More dominance results: incorporating weights and poverty lines

When there are only two variables the conditions are:

## More dominance results: incorporating weights and poverty lines

When there are only two variables the conditions are:

$$\overline{F^A}(x, y) \geq \overline{F^B}(x, y) \forall (x, y) \in [x_{\min}, x_{\max}] \times [y_{\min}, y_{\max}]$$

$$F^A(x, y) \leq F^B(x, y) \forall (x, y) \in [x_{\min}, x_{\max}] \times [y_{\min}, y_{\max}]$$

## More dominance results: incorporating weights and poverty lines

When there are only two variables the conditions are:

$$\overline{F^A}(x, y) \geq \overline{F^B}(x, y) \forall (x, y) \in [x_{\min}, x_{\max}] \times [y_{\min}, y_{\max}]$$

$$F^A(x, y) \leq F^B(x, y) \forall (x, y) \in [x_{\min}, x_{\max}] \times [y_{\min}, y_{\max}]$$

When there are several variables and the poor are identified according to the *union* approach:

## More dominance results: incorporating weights and poverty lines

When there are only two variables the conditions are:

$$\overline{F^A}(x, y) \geq \overline{F^B}(x, y) \forall (x, y) \in [x_{\min}, x_{\max}] \times [y_{\min}, y_{\max}]$$

$$F^A(x, y) \leq F^B(x, y) \forall (x, y) \in [x_{\min}, x_{\max}] \times [y_{\min}, y_{\max}]$$

When there are several variables and the poor are identified according to the *union* approach:

$$M_A^\alpha(X; \min(w_d); Z) \leq M_B^\alpha(X; \min(w_d); Z) \forall \alpha \in \mathbb{R}_0^+$$

$$\forall w_d \in \mathbb{R}^+ \mid \sum_{d=1}^D w_d = 1, \forall Z \leftrightarrow F_d^A(x_d) \leq F_d^B(x_d) \forall x_1, \dots, x_D$$

## More dominance results: incorporating weights and poverty lines

When there are several variables and the poor are identified according to the *intersection* approach:

## More dominance results: incorporating weights and poverty lines

When there are several variables and the poor are identified according to the *intersection* approach:

$$M_A^\alpha(X; D; Z) \leq M_B^\alpha(X; D; Z) \forall \alpha \in \mathbb{R}_0^+$$

$$\forall w_d \in \mathbb{R}^+ \mid \sum_{d=1}^D w_d = 1, \forall Z \leftrightarrow$$

$$F^A(x_1, \dots, x_D) \leq F^B(x_1, \dots, x_D) \forall x_1, \dots, x_D$$

# Illustration of the conditions for $H(X; w, k, Z)$

$$A = \begin{array}{cc} & \begin{matrix} y_1 & y_2 \end{matrix} \\ \begin{matrix} x_1 \\ x_2 \end{matrix} & \begin{bmatrix} 0.3 & 0.1 \\ 0.1 & 0.5 \end{bmatrix} \end{array}$$

$$B = \begin{array}{cc} & \begin{matrix} y_1 & y_2 \end{matrix} \\ \begin{matrix} x_1 \\ x_2 \end{matrix} & \begin{bmatrix} 0.2 & 0.2 \\ 0.2 & 0.4 \end{bmatrix} \end{array}$$

$$C = \begin{array}{cc} & \begin{matrix} y_1 & y_2 \end{matrix} \\ \begin{matrix} x_1 \\ x_2 \end{matrix} & \begin{bmatrix} 0.2 & 0.15 \\ 0.15 & 0.5 \end{bmatrix} \end{array}$$



# Illustration of the conditions for $H(X; w, k, Z)$

Table 1: Multidimensional headcount ratios

Values of $c_n$	$w_x$ vs $w_y$	$z_x, z_y$	$H^A$	$H^B$	$H^C$
$w_x$	$w_x > w_y$	$x_1, y_1$	0.4	0.4	0.35
$w_x$	$w_x \leq w_y$	$x_1, y_1$	0.5	0.6	0.5
$w_x$	$w_x > w_y$	$x_1, y_2$	0.4	0.4	0.35
$w_x$	$w_x \leq w_y$	$x_1, y_2$	1	1	1
$w_x$		$x_2, y_1$	1	1	1
$w_y$	$w_y > w_x$	$x_1, y_1$	0.4	0.4	0.35
$w_y$	$w_y \leq w_x$	$x_1, y_1$	0.5	0.6	0.5
$w_y$		$x_1, y_2$	1	1	1
$w_y$	$w_y > w_x$	$x_2, y_1$	0.4	0.4	0.35
$w_y$	$w_y \leq w_x$	$x_2, y_1$	1	1	1
$w_x + w_y$		$x_1, y_1$	0.3	0.2	0.2
$w_x + w_y$		$x_1, y_2$	0.4	0.4	0.35
$w_x + w_y$		$x_2, y_1$	0.4	0.4	0.35

## Illustration of the conditions for $H(X; w, k, Z)$ : the cumulative and survival distributions

$$\begin{array}{l}
 F^A = \begin{array}{cc} & y_1 & y_2 \\ x_1 & 0.3 & 0.4 \\ x_2 & 0.4 & 1 \end{array} ; F^B = \begin{array}{cc} & y_1 & y_2 \\ x_1 & 0.2 & 0.4 \\ x_2 & 0.4 & 1 \end{array} ; F^C = \begin{array}{cc} & y_1 & y_2 \\ x_1 & 0.2 & 0.35 \\ x_2 & 0.35 & 1 \end{array} \\
 \bar{F}^A = \begin{array}{cc} & y_1 & y_2 \\ x_1 & 1 & 0.6 \\ x_2 & 0.6 & 0.5 \end{array} ; \bar{F}^B = \begin{array}{cc} & y_1 & y_2 \\ x_1 & 1 & 0.6 \\ x_2 & 0.6 & 0.4 \end{array} ; \bar{F}^C = \begin{array}{cc} & y_1 & y_2 \\ x_1 & 1 & 0.65 \\ x_2 & 0.65 & 0.5 \end{array}
 \end{array}$$

## Why do we need other methods?

- ▶ Stochastic dominance conditions are useful for evaluating *ordinal* pairwise comparisons.

## Why do we need other methods?

- ▶ Stochastic dominance conditions are useful for evaluating *ordinal* pairwise comparisons.
- ▶ However, these are very stringent conditions, they are not always fulfilled and they are not (yet) available for many interesting cases.

## Why do we need other methods?

- ▶ Stochastic dominance conditions are useful for evaluating *ordinal* pairwise comparisons.
- ▶ However, these are very stringent conditions, they are not always fulfilled and they are not (yet) available for many interesting cases.
- ▶ What can we say about the degree of robustness of a whole ranking of countries, people, etc. when dominance conditions are not fulfilled for all possible pairs?

## Why do we need other methods?

- ▶ Stochastic dominance conditions are useful for evaluating *ordinal* pairwise comparisons.
- ▶ However, these are very stringent conditions, they are not always fulfilled and they are not (yet) available for many interesting cases.
- ▶ What can we say about the degree of robustness of a whole ranking of countries, people, etc. when dominance conditions are not fulfilled for all possible pairs?
- ▶ We need other methods to quantify and assess the robustness of the orderings to changes in the parameters.

## Other methods

In this presentation we will review some of the methods used in the report by Alkire et al. (2010):

## Other methods

In this presentation we will review some of the methods used in the report by Alkire et al. (2010):

- ▶ Correlation coefficients for pairs of rankings, e.g. the Gamma by Goodman and Kruskal, and Spearman's Rho.



## Other methods

In this presentation we will review some of the methods used in the report by Alkire et al. (2010):

- ▶ Correlation coefficients for pairs of rankings, e.g. the Gamma by Goodman and Kruskal, and Spearman's Rho.
- ▶ The multiple rank concordance indices, e.g. Kendall-Friedman, Kendall, Joe.

## Other methods

In this presentation we will review some of the methods used in the report by Alkire et al. (2010):

- ▶ Correlation coefficients for pairs of rankings, e.g. the Gamma by Goodman and Kruskal, and Spearman's Rho.
- ▶ The multiple rank concordance indices, e.g. Kendall-Friedman, Kendall, Joe.
- ▶ Percentages of reversed comparisons.

## Other methods

In this presentation we will review some of the methods used in the report by Alkire et al. (2010):

- ▶ Correlation coefficients for pairs of rankings, e.g. the Gamma by Goodman and Kruskal, and Spearman's Rho.
- ▶ The multiple rank concordance indices, e.g. Kendall-Friedman, Kendall, Joe.
- ▶ Percentages of reversed comparisons.
- ▶ Counting of large changes in the positions.

## The context

- ▶ There are  $N$  countries

## The context

- ▶ There are  $N$  countries
- ▶ The ordering/ranking of the  $N$  countries is represented by the vector:

$$R = (R_1, R_2, \dots, R_N)$$

We assume:  $R_1 < R_2 < \dots < R_N$ .

## The context

- ▶ There are  $N$  countries
- ▶ The ordering/ranking of the  $N$  countries is represented by the vector:

$$R = (R_1, R_2, \dots, R_N)$$

We assume:  $R_1 < R_2 < \dots < R_N$ .

- ▶ The objective is to analyze the robustness of ranking  $R$  for alternative parameter values.

## The context

- ▶ There are  $N$  countries
- ▶ The ordering/ranking of the  $N$  countries is represented by the vector:

$$R = (R_1, R_2, \dots, R_N)$$

We assume:  $R_1 < R_2 < \dots < R_N$ .

- ▶ The objective is to analyze the robustness of ranking  $R$  for alternative parameter values.
- ▶ Maybe the changes are in the weights,  $w'$ ; poverty lines,  $z'$ ; different thresholds,  $k'$ ; even the number of dimensions might change to  $D'$ .

# The context



## The context

- ▶ The new ranking under a different specification is denoted by the vector:

$$R' = (R'_1, R'_2, \dots, R'_N)$$

## The context

- ▶ The new ranking under a different specification is denoted by the vector:

$$R' = (R'_1, R'_2, \dots, R'_N)$$

- ▶ If  $R_n = R'_n$  for all  $n = 1, \dots, N$  then the ordering is completely robust with respect to the alternative specification.

## The context

- ▶ The new ranking under a different specification is denoted by the vector:

$$R' = (R'_1, R'_2, \dots, R'_N)$$

- ▶ If  $R_n = R'_n$  for all  $n = 1, \dots, N$  then the ordering is completely robust with respect to the alternative specification.
- ▶ In general, we want to measure how close two, or more, rankings, stand from a situation of perfect correlation, or concordance, and which classified elements may be responsible for any deviation from complete robustness.

## Rank correlations

- ▶ A useful method is to measure the correlation between different orderings.

## Rank correlations

- ▶ A useful method is to measure the correlation between different orderings.
- ▶ We review two methods:

## Rank correlations

- ▶ A useful method is to measure the correlation between different orderings.
- ▶ We review two methods:
  - ▶ The rank correlation coefficient of Goodman y Kruskal:  $(\gamma)$ .

## Rank correlations

- ▶ A useful method is to measure the correlation between different orderings.
- ▶ We review two methods:
  - ▶ The rank correlation coefficient of Goodman y Kruskal:  $(\gamma)$ .
  - ▶ The rank correlation coefficient of Spearman:  $(\rho)$ .

## Rank correlations

- ▶ A useful method is to measure the correlation between different orderings.
- ▶ We review two methods:
  - ▶ The rank correlation coefficient of Goodman y Kruskal:  $(\gamma)$ .
  - ▶ The rank correlation coefficient of Spearman:  $(\rho)$ .



## The $\gamma$ of Goodman and Kruskal

- ▶  $\gamma$  is based on:

## The $\gamma$ of Goodman and Kruskal

- ▶  $\gamma$  is based on:
  - ▶ The number of concordant pairs ( $C$ )

## The $\gamma$ of Goodman and Kruskal

- ▶  $\gamma$  is based on:
  - ▶ The number of concordant pairs ( $C$ )
  - ▶ The number of discordant pairs ( $D$ )

## The $\gamma$ of Goodman and Kruskal

- ▶  $\gamma$  is based on:
  - ▶ The number of concordant pairs ( $C$ )
  - ▶ The number of discordant pairs ( $D$ )
- ▶ Concordant and discordant pairs:
  - ▶ A pair  $n$  and  $\bar{n}$  is concordant if:

$$R_n > R_{\bar{n}} \text{ and } R'_n > R'_{\bar{n}}$$

## The $\gamma$ of Goodman and Kruskal

- ▶  $\gamma$  is based on:
  - ▶ The number of concordant pairs ( $C$ )
  - ▶ The number of discordant pairs ( $D$ )
- ▶ Concordant and discordant pairs:
  - ▶ A pair  $n$  and  $\bar{n}$  is concordant if:

$$R_n > R_{\bar{n}} \text{ and } R'_n > R'_{\bar{n}}$$

- ▶ The pair is discordant if:

$$R_n > R_{\bar{n}} \text{ but } R'_n < R'_{\bar{n}}$$

## The $\gamma$ of Goodman and Kruskal

- ▶  $\gamma$  is based on:
  - ▶ The number of concordant pairs ( $C$ )
  - ▶ The number of discordant pairs ( $D$ )
- ▶ Concordant and discordant pairs:
  - ▶ A pair  $n$  and  $\bar{n}$  is concordant if:

$$R_n > R_{\bar{n}} \text{ and } R'_n > R'_{\bar{n}}$$

- ▶ The pair is discordant if:

$$R_n > R_{\bar{n}} \text{ but } R'_n < R'_{\bar{n}}$$

- ▶ With these concepts,  $\gamma$  is defined as:

$$\gamma = \frac{C - D}{C + D}$$

## Exploring Gamma

- ▶ Gamma measures the normalized difference between the total of concordant pairs and discordant pairs.

## Exploring Gamma

- ▶ Gamma measures the normalized difference between the total of concordant pairs and discordant pairs.
- ▶ Notice that  $C + D$  is the total number of comparisons when there are no ties among the ranks.



## Exploring Gamma

- ▶ Gamma measures the normalized difference between the total of concordant pairs and discordant pairs.
- ▶ Notice that  $C + D$  is the total number of comparisons when there are no ties among the ranks. Since  $M_0$  is continuous, we assume that there are no ties.

## Exploring Gamma

- ▶ Gamma measures the normalized difference between the total of concordant pairs and discordant pairs.
- ▶ Notice that  $C + D$  is the total number of comparisons when there are no ties among the ranks. Since  $M_0$  is continuous, we assume that there are no ties.
- ▶ The maximum value of  $\gamma$  is  $+1$  (perfect rank correlation).

## Exploring Gamma

- ▶ Gamma measures the normalized difference between the total of concordant pairs and discordant pairs.
- ▶ Notice that  $C + D$  is the total number of comparisons when there are no ties among the ranks. Since  $M_0$  is continuous, we assume that there are no ties.
- ▶ The maximum value of  $\gamma$  is  $+1$  (perfect rank correlation).  
 $R_n = R'_n$  for all  $n$ .
- ▶ The minimum value of  $\gamma$  is  $-1$  (all pairs are discordant).

## Exploring Gamma

- ▶ Gamma measures the normalized difference between the total of concordant pairs and discordant pairs.
- ▶ Notice that  $C + D$  is the total number of comparisons when there are no ties among the ranks. Since  $M_0$  is continuous, we assume that there are no ties.
- ▶ The maximum value of  $\gamma$  is  $+1$  (perfect rank correlation).  
 $R_n = R'_n$  for all  $n$ .
- ▶ The minimum value of  $\gamma$  is  $-1$  (all pairs are discordant). One ranking is the exact reverse of the other one.

## Exploring Gamma

- ▶ Gamma measures the normalized difference between the total of concordant pairs and discordant pairs.
- ▶ Notice that  $C + D$  is the total number of comparisons when there are no ties among the ranks. Since  $M_0$  is continuous, we assume that there are no ties.
- ▶ The maximum value of  $\gamma$  is  $+1$  (perfect rank correlation).  
 $R_n = R'_n$  for all  $n$ .
- ▶ The minimum value of  $\gamma$  is  $-1$  (all pairs are discordant). One ranking is the exact reverse of the other one.
- ▶ When the number of concordant pairs is equal to that of discordant pairs:  $\gamma = 0$ .

## Spearman's correlation coefficient

- ▶ Rho is based on the differences in rankings for every classified object (e.g. a country) under two specifications.

## Spearman's correlation coefficient

- ▶ Rho is based on the differences in rankings for every classified object (e.g. a country) under two specifications.
- ▶ We define  $r_n = R_n - R'_n$  for all  $n = 1, \dots, N$ .

## Spearman's correlation coefficient

- ▶ Rho is based on the differences in rankings for every classified object (e.g. a country) under two specifications.
- ▶ We define  $r_n = R_n - R'_n$  for all  $n = 1, \dots, N$ .
  - ▶  $r_n$  is the difference in rankings for country  $n$  under two different ranking criteria.



## Spearman's correlation coefficient

- ▶ Rho is based on the differences in rankings for every classified object (e.g. a country) under two specifications.
- ▶ We define  $r_n = R_n - R'_n$  for all  $n = 1, \dots, N$ .
  - ▶  $r_n$  is the difference in rankings for country  $n$  under two different ranking criteria.
- ▶ Thus Rho has the following expression:

$$\rho = 1 - \frac{6 \sum_{n=1}^N r_n^2}{N(N^2 - 1)}$$

## Spearman's correlation coefficient

- ▶ Rho is based on the differences in rankings for every classified object (e.g. a country) under two specifications.
- ▶ We define  $r_n = R_n - R'_n$  for all  $n = 1, \dots, N$ .
  - ▶  $r_n$  is the difference in rankings for country  $n$  under two different ranking criteria.
- ▶ Thus Rho has the following expression:

$$\rho = 1 - \frac{6 \sum_{n=1}^N r_n^2}{N(N^2 - 1)}$$

- ▶ When  $R_n = R'_n$ :  $r_n = 0$  for all  $n$  and, in turn,  $\rho = +1$ .

## Spearman's correlation coefficient

- ▶ Rho is based on the differences in rankings for every classified object (e.g. a country) under two specifications.
- ▶ We define  $r_n = R_n - R'_n$  for all  $n = 1, \dots, N$ .
  - ▶  $r_n$  is the difference in rankings for country  $n$  under two different ranking criteria.
- ▶ Thus Rho has the following expression:

$$\rho = 1 - \frac{6 \sum_{n=1}^N r_n^2}{N(N^2 - 1)}$$

- ▶ When  $R_n = R'_n$ :  $r_n = 0$  for all  $n$  and, in turn,  $\rho = +1$ .
- ▶ When  $R_n = R'_{N-n+1}$  for all  $n$ :  $\rho = -1$ .

# Illustration: Robustness of weights of the MPI

Table 5: Correlations among MPIs adjusted by Survey

Pair of Rankings Compared	Correlation Coefficient	All Countries	Bottom 75 Countries	DHS Only	MICS Only	DHS & MICS	Non WHS	WHS Only
MPI with 50% weight Education and MPI with 50% weight on Health	Spearman	0.957	0.970	0.980	0.949	0.981	0.979	0.616
	Gamma	0.836	0.854	0.885	0.842	0.883	0.880	0.472
MPI with 50% weight Education and MPI with 50% weight on Living Stan.	Spearman	0.970	0.951	0.946	0.971	0.970	0.972	0.859
	Gamma	0.854	0.805	0.819	0.872	0.856	0.860	0.715
MPI with 50% weight Health and MPI with 50% weight on Living Stan.	Spearman	0.968	0.968	0.965	0.975	0.980	0.978	0.763
	Gamma	0.856	0.854	0.849	0.882	0.881	0.876	0.639
Total Number of Observations		104	75	48	35	83	85	19

## Concordance indices for multiple rankings

Unlike correlation coefficients, these indices provide an evaluation of the degree of concordance of several rankings at the same time.

## Concordance indices for multiple rankings

Unlike correlation coefficients, these indices provide an evaluation of the degree of concordance of several rankings at the same time.

We review three indices for situations where there are not ties (for adjustments for ties see Seth and Yalonetzky, 2011):

## Concordance indices for multiple rankings

Unlike correlation coefficients, these indices provide an evaluation of the degree of concordance of several rankings at the same time.

We review three indices for situations where there are not ties (for adjustments for ties see Seth and Yalonetzky, 2011):

- The index by Kendall-Friedman:

$$KF = \frac{12}{(N^2 - 1)ND^2} \sum_{n=1}^N \left[ \sum_{j=1}^J R_n^j - \frac{(N+1)D}{2} \right]^2$$

## Concordance indices for multiple rankings

Unlike correlation coefficients, these indices provide an evaluation of the degree of concordance of several rankings at the same time.

We review three indices for situations where there are not ties (for adjustments for ties see Seth and Yalonetzky, 2011):

- The index by Kendall-Friedman:

$$KF = \frac{12}{(N^2 - 1)ND^2} \sum_{n=1}^N \left[ \sum_{j=1}^J R_n^j - \frac{(N+1)D}{2} \right]^2$$

- The index by Kendall:

$$K = \frac{12}{N^2 - 1} \left[ \frac{2}{ND(D-1)} \sum_{j=1}^J \sum_{j' > j}^J \sum_{n=1}^N R_n^j R_n^{j'} - \frac{(N+1)^2}{4} \right]$$



## Concordance indices for multiple rankings

Unlike correlation coefficients, these indices provide an evaluation of the degree of concordance of several rankings at the same time.

We review three indices for situations where there are not ties (for adjustments for ties see Seth and Yalonetzky, 2011):

- The index by Kendall-Friedman:

$$KF = \frac{12}{(N^2 - 1)ND^2} \sum_{n=1}^N \left[ \sum_{j=1}^J R_n^j - \frac{(N+1)D}{2} \right]^2$$

- The index by Kendall:

$$K = \frac{12}{N^2 - 1} \left[ \frac{2}{ND(D-1)} \sum_{j=1}^J \sum_{j' > j}^J \sum_{n=1}^N R_n^j R_n^{j'} - \frac{(N+1)^2}{4} \right]$$

- The index by Joe:

$$JO = \frac{\frac{1}{N} \sum_{n=1}^N \prod_{j=1}^J R_n^j - \left(\frac{N+1}{2}\right)^2}{\frac{1}{N} \sum_{n=1}^N n^J - \left(\frac{N+1}{2}\right)^2}$$

## Illustration: Robustness of weights of MPI

Table 6: Indicators of rank concordance for all Survey Sources

Index	All Countries	Bottom 75 Countries	DHS Only	MICS Only	DHS & MICS	Non WHS	WHS Only
KF	0.981	0.981	0.981	0.980	0.987	0.987	0.863
K	0.975	0.974	0.975	0.973	0.983	0.983	0.817
J	0.983	0.974	0.968	0.984	0.982	0.983	0.862

## Other methods: percentage of reversed pairwise comparisons

If  $q$  is the number of pairwise comparisons that, under some criterion, gets reversed (several criteria can be considered simultaneously):

## Other methods: percentage of reversed pairwise comparisons

If  $q$  is the number of pairwise comparisons that, under some criterion, gets reversed (several criteria can be considered simultaneously):

$$0 \leq q \leq \frac{N(N-1)}{2}$$

## Other methods: percentage of reversed pairwise comparisons

If  $q$  is the number of pairwise comparisons that, under some criterion, gets reversed (several criteria can be considered simultaneously):

$$0 \leq q \leq \frac{N(N-1)}{2}$$

Then one can compute:  $p = \frac{2q}{N(N-1)}$

## Other methods: percentage of reversed pairwise comparisons

If  $q$  is the number of pairwise comparisons that, under some criterion, gets reversed (several criteria can be considered simultaneously):

$$0 \leq q \leq \frac{N(N-1)}{2}$$

Then one can compute:  $p = \frac{2q}{N(N-1)}$

Notice that:

## Other methods: percentage of reversed pairwise comparisons

If  $q$  is the number of pairwise comparisons that, under some criterion, gets reversed (several criteria can be considered simultaneously):

$$0 \leq q \leq \frac{N(N-1)}{2}$$

Then one can compute:  $p = \frac{2q}{N(N-1)}$

Notice that:

- ▶ When  $p = 1$  and  $J = 2$  rankings get reversed completely ( $R_n = R'_{N-n+1}$ ).

## Other methods: percentage of reversed pairwise comparisons

If  $q$  is the number of pairwise comparisons that, under some criterion, gets reversed (several criteria can be considered simultaneously):

$$0 \leq q \leq \frac{N(N-1)}{2}$$

Then one can compute:  $p = \frac{2q}{N(N-1)}$

Notice that:

- ▶ When  $p = 1$  and  $J = 2$  rankings get reversed completely ( $R_n = R'_{N-n+1}$ ).
- ▶ The maximum effect generated by just one country occurs when it moves from one extreme of the ranks to the other.



## Other methods: percentage of reversed pairwise comparisons

If  $q$  is the number of pairwise comparisons that, under some criterion, gets reversed (several criteria can be considered simultaneously):

$$0 \leq q \leq \frac{N(N-1)}{2}$$

Then one can compute:  $p = \frac{2q}{N(N-1)}$

Notice that:

- ▶ When  $p = 1$  and  $J = 2$  rankings get reversed completely ( $R_n = R'_{N-n+1}$ ).
- ▶ The maximum effect generated by just one country occurs when it moves from one extreme of the ranks to the other. If that is the only change then:  $p = \frac{2}{N}$ .