

Time decompositions of the adjusted headcount ratio

Gastón Yalonetzky

Oxford Poverty and Human Development Initiative

OPHI-HDCA Summer School, Delft, 24 August - 3 September 2011.

We are grateful to the World Bank, two anonymous donors and OPHI for financial support

Table of contents

Introduction

Methodology

Basic decompositions

Data

Results

Basic decomposition of MPI

Decomposition of H

Decomposition of A

More general results

Data

Results

Concluding remarks

Introduction

- ▶ The adjusted headcount ratio, M_0 , has nice decomposability properties, e.g. by regions, groups, etc.

Introduction

- ▶ The adjusted headcount ratio, M_0 , has nice decomposability properties, e.g. by regions, groups, etc.
- ▶ In this class we will explore the nice decomposability properties of $\Delta\%M_0$ and its components: $\Delta\%H$ and $\Delta\%A$.

Introduction

- ▶ The adjusted headcount ratio, M_0 , has nice decomposability properties, e.g. by regions, groups, etc.
- ▶ In this class we will explore the nice decomposability properties of $\Delta\%M_0$ and its components: $\Delta\%H$ and $\Delta\%A$.
- ▶ This material is based on Apablaza, Ocampo and Yalonetzky (2010), and on Apablaza and Yalonetzky (2011).

Introduction

- ▶ We are going to study the most basic time decompositions first, i.e. for cross-sections.

Introduction

- ▶ We are going to study the most basic time decompositions first, i.e. for cross-sections.
- ▶ Then we will show some results from Apablaza, Ocampo and Yalonetzky (2010).

Introduction

- ▶ We are going to study the most basic time decompositions first, i.e. for cross-sections.
- ▶ Then we will show some results from Apablaza, Ocampo and Yalonetzky (2010).
- ▶ Then we will discuss more decompositions relevant to panel datasets.

Introduction

- ▶ We are going to study the most basic time decompositions first, i.e. for cross-sections.
- ▶ Then we will show some results from Apablaza, Ocampo and Yalonetzky (2010).
- ▶ Then we will discuss more decompositions relevant to panel datasets.
- ▶ And show some results from Apablaza and Yalonetzky (2011).

Introduction

- ▶ We are going to study the most basic time decompositions first, i.e. for cross-sections.
- ▶ Then we will show some results from Apablaza, Ocampo and Yalonetzky (2010).
- ▶ Then we will discuss more decompositions relevant to panel datasets.
- ▶ And show some results from Apablaza and Yalonetzky (2011).
- ▶ We will finish with some remarks on comparability from Apablaza, Ocampo and Yalonetzky (2010).

Basic Notation

The weighted sum of deprivations

Basic Notation

The weighted sum of deprivations

$$c_n = \sum_{d=1}^D w_d I(x_{nd}^t \leq z_d)$$

Basic Notation

The weighted sum of deprivations

$$c_n = \sum_{d=1}^D w_d I(x_{nd}^t \leq z_d)$$

Multidimensional poverty headcount:

$$H(X^t; Z) \equiv \frac{1}{N^t} \sum_{n=1}^{N^t} I(c_n \geq k)$$

Basic Notation

Average deprivation of the poor:

Basic Notation

Average deprivation of the poor:

$$A(X^t; Z) \equiv \frac{1}{DN^t H} \sum_{n=1}^{N^t} I(c_n \geq k) c_n$$

Basic Notation

Average deprivation of the poor:

$$A(X^t; Z) \equiv \frac{1}{DN^t H} \sum_{n=1}^{N^t} I(c_n \geq k) c_n$$

The adjusted headcount ratio, M^0 :

$$M^0(X^t; Z) \equiv H^t A^t = \frac{1}{DN^t} \sum_{n=1}^{N^t} I(c_n \geq k) c_n$$

Basic Notation

Average deprivation of the poor:

$$A(X^t; Z) \equiv \frac{1}{DN^t H} \sum_{n=1}^{N^t} I(c_n \geq k) c_n$$

The adjusted headcount ratio, M^0 :

$$M^0(X^t; Z) \equiv H^t A^t = \frac{1}{DN^t} \sum_{n=1}^{N^t} I(c_n \geq k) c_n$$

$$\Delta\%_a M^0(t) \equiv \frac{M^0(X^t; Z) - M^0(X^{t-a}; Z)}{M^0(X^{t-a}; Z)}$$

Basic decomposition of M^0

$$\Delta\%_a M^0(t) = \Delta\%_a H(t) + \Delta\%_a A(t) + \Delta\%_a H(t)\Delta\%_a A(t)$$

- ▶ $\Delta\%_a H(t)$ and $\Delta\%_a A(t)$ are not generally independent, but sometimes a change in one may not produce a change in the other.

Basic decomposition of M^0

$$\Delta\%_a M^0(t) = \Delta\%_a H(t) + \Delta\%_a A(t) + \Delta\%_a H(t)\Delta\%_a A(t)$$

- ▶ $\Delta\%_a H(t)$ and $\Delta\%_a A(t)$ are not generally independent, but sometimes a change in one may not produce a change in the other.
- ▶ E.g. if $k = D$: $\Delta\%_a A(t) = 0$, and $\Delta\%_a M(t) = \Delta\%_a H(t)$.

Basic decomposition of M^0

$$\Delta\%_a M^0(t) = \Delta\%_a H(t) + \Delta\%_a A(t) + \Delta\%_a H(t)\Delta\%_a A(t)$$

- ▶ $\Delta\%_a H(t)$ and $\Delta\%_a A(t)$ are not generally independent, but sometimes a change in one may not produce a change in the other.
- ▶ E.g. if $k = D$: $\Delta\%_a A(t) = 0$, and $\Delta\%_a M(t) = \Delta\%_a H(t)$.
- ▶ or if $k < D$ it is possible that $\Delta\%_a H(t) = 0$ and $\Delta\%_a A(t) \neq 0$.

Basic decomposition of M^0

$$\Delta\%_a M^0(t) = \Delta\%_a H(t) + \Delta\%_a A(t) + \Delta\%_a H(t)\Delta\%_a A(t)$$

- ▶ $\Delta\%_a H(t)$ and $\Delta\%_a A(t)$ are not generally independent, but sometimes a change in one may not produce a change in the other.
- ▶ E.g. if $k = D$: $\Delta\%_a A(t) = 0$, and $\Delta\%_a M(t) = \Delta\%_a H(t)$.
- ▶ or if $k < D$ it is possible that $\Delta\%_a H(t) = 0$ and $\Delta\%_a A(t) \neq 0$.
- ▶ As k goes from 1 to D , H decreases and A increases "mechanically". Hence as k increases toward D , it is more likely to find higher $\Delta\%_a H(t)$ and lower $\Delta\%_a A(t)$.

Basic decomposition of H

H is decomposable into the multidimensional headcounts of subgroups:

$$H(X^t, Z) = \sum_{i=1}^G \psi_i^t H^i(X_i^t, Z)$$

Basic decomposition of H

H is decomposable into the multidimensional headcounts of subgroups:

$$H(X^t, Z) = \sum_{i=1}^G \psi_i^t H^i(X_i^t, Z)$$

Where $\psi_i^t \equiv \frac{N_i^t}{N^t}$

Basic decomposition of H

H is decomposable into the multidimensional headcounts of subgroups:

$$H(X^t, Z) = \sum_{i=1}^G \psi_i^t H^i(X_i^t, Z)$$

Where $\psi_i^t \equiv \frac{N_i^t}{N^t}$

In turn:

$$H^i(X_i^t, Z) = \frac{1}{N_i^t} \sum_{n=1}^N I(c_n \geq k) I(n \in i)$$

Basic decomposition of H

$$\Delta\%_a H(t) = \sum_{i=1}^G \Delta\%_a [\psi_i^t H^i(X_i^t, Z)] r_i(t-a)$$

Basic decomposition of H

$$\Delta\%_a H(t) = \sum_{i=1}^G \Delta\%_a [\psi_i^t H^i(X_i^t, Z)] r_i(t-a)$$

Where $r_i(t-a) \equiv \frac{\psi_i^{t-a} H^i(X_i^{t-a}, Z)}{H(X^{t-a}, Z)}$

Basic decomposition of H

$$\Delta\%_a H(t) = \sum_{i=1}^G \Delta\%_a [\psi_i^t H^i(X_i^t, Z)] r_i(t-a)$$

Where $r_i(t-a) \equiv \frac{\psi_i^{t-a} H^i(X_i^{t-a}, Z)}{H(X^{t-a}, Z)}$

Then:

$$\Delta\%_a H(t) = \sum_{i=1}^G r_i(t-a) [\Delta\%_a \psi_i^t + \Delta\%_a H^i(X_i^t, Z) + \Delta\%_a \psi_i^t \Delta\%_a H^i(X_i^t, Z)]$$

Basic decomposition of A

$$\Delta\%_a A(t) = \sum_{d=1}^D \Delta\%_a [\theta_d A_d(X^t, Z)] s_d(t-a)$$

Basic decomposition of A

$$\Delta\%_a A(t) = \sum_{d=1}^D \Delta\%_a [\theta_d A_d(X^t, Z)] s_d(t-a)$$

Where $\theta_d \equiv \frac{w_d}{D}$ and $s_d(t-a) \equiv \frac{\theta_d A_d(X^{t-a}, Z)}{A(X^{t-a}, Z)}$

Basic decomposition of A

$$\Delta\%_a A(t) = \sum_{d=1}^D \Delta\%_a [\theta_d A_d(X^t, Z)] s_d(t-a)$$

Where $\theta_d \equiv \frac{w_d}{D}$ and $s_d(t-a) \equiv \frac{\theta_d A_d(X^{t-a}, Z)}{A(X^{t-a}, Z)}$

And $A_d(X^t, Z) \equiv \frac{\sum_{n=1}^{N^t} I(c_n \geq k \wedge x_{nd}^t \leq z_d)}{N(t)H(t)}$

Then:

$$\Delta\%_a A(t) = \sum_{d=1}^D s_d(t-a)[\Delta\%_a \theta_d A_d(X^t, Z)] = \sum_{d=1}^D s_d(t-a)[\Delta\%_a A_d(X^t, Z)]$$

Then:

$$\Delta\%_a A(t) = \sum_{d=1}^D s_d(t-a) [\Delta\%_a \theta_d A_d(X^t, Z)] = \sum_{d=1}^D s_d(t-a) [\Delta\%_a A_d(X^t, Z)]$$

Because, by construction, $\Delta\%_a \theta_d = 0$

Then:

$$\Delta\%_a A(t) = \sum_{d=1}^D s_d(t-a) [\Delta\%_a \theta_d A_d(X^t, Z)] = \sum_{d=1}^D s_d(t-a) [\Delta\%_a A_d(X^t, Z)]$$

Because, by construction, $\Delta\%_a \theta_d = 0$

In practical comparisons, we divide the changes by the year-gaps to improve comparability.

The countries

Country	Years
Bangladesh	2004-2007
Colombia	1995-2005
Ethiopia	2000-2005
Ghana	2003-2008
India	1999-2005
Morocco	1992-2004
Nepal	2001-2006
Nigeria	1999-2003
Tanzania	2005-2008
Vietnam	1997-2002


The variables

Variable	B	C	E	G	I	M	Ne	Ni	T	V
Years school	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Enrollment	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Child mortality	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Nutrition	✓	✓	✓	✓	✓	✓	✓	✓	x	x
Electricity	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Toilet	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Water	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Floor	✓	✓	✓	✓	x	✓	✓	✓	✓	✓
Cooking	✓	✓	✓	✓	✓	x	✓	x	✓	x
Asset	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓


B=Bangladesh; C=Colombia; E=Ethiopia; G=Ghana; I=India

M=Morocco; Ne=Nepal; Ni=Nigeria; T=Tanzania; V=Vietnam


Decomposition of M0 for 10 countries and k=3




The impact of the choice of k: the case of Bangladesh




The impact of the choice of k: the case of Colombia




The impact of the choice of k: the case of Ethiopia




The impact of the choice of k: the case of Ghana




The impact of the choice of k: the case of India




The impact of the choice of k: the case of Morocco




The impact of the choice of k: the case of Nepal




The impact of the choice of k: the case of Nigeria




The impact of the choice of k: the case of Tanzania




The impact of the choice of k: the case of Vietnam



Decomposition of H for k=3



Decomposition of A for k=3



More general results for $\Delta\%_a M0$

Consider now the censored headcount, $CH_d(t)$:

$$CH_d(t) \equiv \frac{1}{N^t} \sum_{n=1}^{N^t} I(c_n \geq k \wedge x_{nd}^t \leq z_d)$$

More general results for $\Delta\%_a M^0$

Consider now the censored headcount, $CH_d(t)$:

$$CH_d(t) \equiv \frac{1}{N^t} \sum_{n=1}^{N^t} I(c_n \geq k \wedge x_{nd}^t \leq z_d)$$

Then: $A_d = \frac{CH_d(t)}{H(t)}$ and $M^0(t) = \sum_{d=1}^D \theta_d CH_d(t)$

More general results for $\Delta\%_a M0$

We get additional results that rely on $\Delta\%_a CH_d(t)$:

More general results for $\Delta\%_a M_0$

We get additional results that rely on $\Delta\%_a CH_d(t)$:

$$\Delta\%_a A_d = \frac{1 + \Delta\%_a CH_d}{1 + \Delta\%_a H(t)} - 1$$

More general results for $\Delta\%_a M0$

We get additional results that rely on $\Delta\%_a CH_d(t)$:

$$\Delta\%_a A_d = \frac{1 + \Delta\%_a CH_d}{1 + \Delta\%_a H(t)} - 1$$

$$\Delta\%_a A = \frac{1 + \sum_{d=1}^D s_d(t-a) \Delta\%_a CH_d}{1 + \Delta\%_a H(t)} - 1$$

More general results for $\Delta\%_a M^0$

We get additional results that rely on $\Delta\%_a CH_d(t)$:

$$\Delta\%_a A_d = \frac{1 + \Delta\%_a CH_d}{1 + \Delta\%_a H(t)} - 1$$

$$\Delta\%_a A = \frac{1 + \sum_{d=1}^D s_d(t-a) \Delta\%_a CH_d}{1 + \Delta\%_a H(t)} - 1$$

$$\Delta\%_a M^0(t) = \sum_{d=1}^D s_d(t-a) \Delta\%_a CH_d$$

Linking changes to transition probabilities

Now we can link this chain of changes to changes in the transition probabilities of H and CH .

Linking changes to transition probabilities

Now we can link this chain of changes to changes in the transition probabilities of H and CH .

First we have the two "laws of motion":

Linking changes to transition probabilities

Now we can link this chain of changes to changes in the transition probabilities of H and CH .

First we have the two "laws of motion":

$$\Delta^0_a H(t) = P[c_n^t \geq k | c_n^{t-a} < k] \left[\frac{1 - H(t-a)}{H(t-a)} \right] - P[c_n^t < k | c_n^{t-a} \geq k]$$

Linking changes to transition probabilities


Now we can link this chain of changes to changes in the transition probabilities of H and CH .

First we have the two "laws of motion":

$$\Delta\%_a H(t) = P[c_n^t \geq k | c_n^{t-a} < k] \left[\frac{1 - H(t-a)}{H(t-a)} \right] - P[c_n^t < k | c_n^{t-a} \geq k]$$


$$\begin{aligned} \Delta\%_a CH(t) = & P[c_n^t \geq k \wedge x_{nd}^t \leq z_d | c_n^{t-a} < k \vee x_{nd}^{t-a} > z_d] \left[\frac{1 - CH(t-a)}{CH(t-a)} \right] \\ & - P[c_n^t < k \vee x_{nd}^t > z_d | c_n^{t-a} \geq k \wedge x_{nd}^{t-a} \leq z_d] \end{aligned}$$

Decomposition of Alkire-Foster statistics based on transition probabilities




Two final results

I:



Two final results

II:



The Young Lives dataset

We use the three waves: 2002, 2006/7, 2010.

Table 1: Sample Characteristics


	Wave	Original sample	Selected Sample	Mean Age	% Females	% rural
Ethiopia	1	1000	868	7.88	49.1%	61.2%
	2	980	868	12.05		60.7%
	3	973	868	14.56		59.7%
Andhra Pradesh	1	1008	944	7.98	50.6%	75.6%
	2	994	944	12.32		74.8%
	3	975	944	14.72		57.1%
Peru	1	714	660	7.93	47.0%	26.1%
	2	685	660	12.31		40.3%
	3	678	660	14.44		23.6%
Vietnam	1	1000	957	7.97	50.4%	80.6%
	2	990	957	12.25		69.3%
	3	974	957	14.73		n.a.


Choice of variables

Table 2: Child Poverty Dimensions

Indicator	Description (threshold)	Weight
Child Related		
Child Labour [♣]	Any "commercial" activity before 13 / Light activity from 13 (2 hours per day)	1/12%
School Attendance	No attendance to the school according to National Law	1/12%
Attachment	Any contact with parents mum or dad	1/12%
Nutrition [◇]	Less than 2 standards deviations (BMI)	1/12%
Household Related		
Electricity	No electricity	1/12%
Cooking Fuel	MDG definition (Branches/ Charcoal/ Coal/ Cow dung /Crop residues / Leaves/ None /Other)	1/12%
Drinking Water	MDG definition (Unprotected/ Well/ Spring/ Pond/ River/ Stream / Canal)	1/12%
Toilet	MDG definition (Forest/ field/ Open place / Neighbours toilet/ Communal pit latrine/ Relative's toilet/ Simple latrine on pond/ Toilet in health post/ Other)	1/12%
Floor	MDG definition (Earth/ Sand)	1/12%
Assets	Less than one (Radio/ Fridge/ Table/ Bike/ Tv/ Motorbike/ Car/ Phone)	1/12%
Overcrowding [♣]	3 or more Individuals per room	1/12%
Child Mortality [♡]	Any dead Children in the Household	1/12%

Transition probabilities



Transition probabilities and $\Delta\%H$ 

Concluding remarks on time comparisons with M0 across countries

When time periods differ three potential problems of comparability arise:
(Apablaza, Ocampo and Yalonetzky, 2010)

Concluding remarks on time comparisons with M0 across countries

When time periods differ three potential problems of comparability arise:
(Apablaza, Ocampo and Yalonetzky, 2010)

- ▶ Time spans are different: It may give undue advantage for, say, poverty reduction to the country with the longest observed span.
Solution: Annualize rates of change.

Concluding remarks on time comparisons with M0 across countries

When time periods differ three potential problems of comparability arise: (Apablaza, Ocampo and Yalonetzky, 2010)

- ▶ Time spans are different: It may give undue advantage for, say, poverty reduction to the country with the longest observed span. Solution: Annualize rates of change.
- ▶ Time spans are different: If the differences are too wild it may be that for one country we observe short-term business cycle fluctuations whereas for the other we observe a medium-term growth trend. Solution: Restrict comparisons to time spans that do not differ too wildly.

Concluding remarks on time comparisons with M0 across countries

When time periods differ three potential problems of comparability arise: (Apablaza, Ocampo and Yalonetzky, 2010)

- ▶ Time spans are different: It may give undue advantage for, say, poverty reduction to the country with the longest observed span. Solution: Annualize rates of change.
- ▶ Time spans are different: If the differences are too wild it may be that for one country we observe short-term business cycle fluctuations whereas for the other we observe a medium-term growth trend. Solution: Restrict comparisons to time spans that do not differ too wildly.
- ▶ The years are different: Even when spans are equal, taking year brackets too far apart may affect the meaningfulness of the comparison. (E.g. Kenya in the 1950s with Chile in the 1990s). Solution: Justify your comparisons when the years are different.